



Institute of Materia Medica, Chinese Academy of Medical Sciences
Chinese Pharmaceutical Association

Acta Pharmaceutica Sinica B

www.elsevier.com/locate/apsb
www.sciencedirect.com



ORIGINAL ARTICLE

Catalytic cleavage activities of 10–23 DNzyme analogs functionalized with an amino group in its catalytic core

Qi Wang^a, Di Zhang^b, Maosheng Cheng^b, Junlin He^{a,*}, Keliang Liu^{a,*}

^aBeijing Institute of Pharmacology and Toxicology, Beijing 100850, China

^bSchool of Pharmaceutical Engineering, Shenyang Pharmaceutical University, Shenyang 110016, China

Received 30 September 2011; revised 26 October 2011; accepted 29 November 2011

KEY WORDS

10–23 DNzyme;
8-Aza-7-deaza-2'-deoxy-adenosine;
Amino group;
Chemical modification

Abstract Functionalization of the catalytic loop of 10–23 DNzyme with an amino group was performed by incorporation of 7-(3-aminopropyl)-8-aza-7-deaza-2'-deoxyadenosine in different single positions. Among the nine modified positions in the catalytic loop, A9 is the unique position with positive contribution by such modification. These results indicated that more efficient deoxyribozymes remain to be explored by introduction of exogenous functional groups in an appropriate position in the catalytic loop of 10–23 DNzyme, such as the combination of 7-functional group substituted 8-aza-7-deaza-2'-deoxyadenosine analogs and A9 position.

© 2012 Institute of Materia Medica, Chinese Academy of Medical Sciences and Chinese Pharmaceutical Association. Production and hosting by Elsevier B.V. All rights reserved.

*Corresponding authors. Tel.: +6 10 68169363, +6 10 68237761.

E-mail addresses: hejunlin@yahoo.com (Junlin He), keliangliu55@126.com, keliangliu@yahoo.com (Keliang Liu).



1. Introduction

10–23 DNAzyme, an unnatural DNA enzyme obtained by *in vitro* selection from a DNA reservoir, is a sequence-specific RNA-cleaving DNA molecule^{1,2}. Compared with ribozymes, its high chemical and biological stability and convenient synthesis seem to be more in favor of therapeutic and other biotechnological applications. Indeed, many efforts have been made for its applications, including the suppression of specific disease-related genes^{3–16}, nucleic acid structural and functional analysis, and other applications like DNA nanomachine^{17–24}. Improvement of its intrinsic characteristics, such as cleavage site preferences, cleavage efficiencies, and high-level Mg^{2+} -dependence, as well as nuclease stability and efficient delivery for its optimal activities has been the focuses of many researches^{25–34}. Some DNAzymes with protein-like functional groups were selected *in vitro*, which could conduct RNA-cleavage reaction with M^{2+} -independence^{29,34–37}. 10–23 DNAzyme analogs with M^{2+} -independence was obtained, but with a sacrifice of cleavage efficiency. Because an insight into the catalytic mechanism of the DNAzyme is not yet available^{38,39}, new random selections and chemical modifications for more efficient variants are still currently necessary^{40–44}. As we have reported, screening with 7-substituted 8-aza-7-deaza-2'-deoxyadenosine derivatives in the positions of five dA residues (A5, A9, A11, A12, and A15) led to the finding that A9 is the right position for 7-substituted 8-aza-7-deaza-deoxyadenosine analogs, by which N8 atom, 7-substituted phenethyl, 3-hydroxypropyl and 3-aminopropyl were introduced and positive effect on cleavage rate was always observed (Fig. 1)⁴⁵. Especially, replacement of 7-(3-aminopropyl)-8-aza-7-deaza-2'-deoxyadenosine (**1**) at A9 position promised a 12-fold increase of cleavage rate. We postulated that the location of the amino group by this modification contributed to a most favorable catalytic conformation. In this study, the effect of this compound in other four non-dA positions were explored, with the aim to evaluate an appropriate position for the positive role of the extra amino group of compound **1** in the catalytic core of 10–23 DNAzyme.




Figure 1 Left: the complex of 10–23 DNAzyme and its DNA–RNA–DNA substrate. Bold letters represent the RNA residues in the substrate. The arrow denotes the cleavage site on the substrate. Right: compound **1**.

2. Results and discussion

The oligodeoxynucleotides were synthesized with phosphoramidite chemistry. The phosphoramidite of compound **1** was synthesized according to the published procedure⁴⁵. The oligodeoxynucleotides were purified by preparative denaturing 20% polyacrylamide gel electrophoresis (containing 7 M urea) and desalted with SEP-PAK cartridges. Characterization was performed with MALDI-TOF MS (Table 1). In the 10–23 DNAzyme analogs, compound **1** was incorporated into four positions, instead of T8, C10, C13 and G14, namely, **DZ-1-8**, **DZ-1-10**, **DZ-1-13** and **DZ-1-14** were obtained, respectively.

In order to demonstrate whether these modified DNAzymes could bind the substrate as 10–23 DNAzyme, the T_m of the complexes of these modified DNAzymes with the substrate was evaluated under the conditions for the cleavage reaction (50 mM Tris–HCl, 2 mM Mg^{2+})⁴⁵, with the chimeric DNA–RNA–DNA substrate replaced by its corresponding full-DNA substrate D19, 5'-d (AGG TGC AGG ATG GAG AGC A)-3' to avoid any cleavage reactions conducted by DNAzyme itself and nucleases in the environment. All the complexes have a T_m of 51 °C (Table 1), very close to that of the complex of 10–23 DNAzyme and D19 (52 °C). It meant that the chemical modifications on the catalytic core could not result in changes on the thermal stability of the complexes. Furthermore, the temperature of all the cleavage reactions was set at 37 °C, much lower than the T_m of DNAzyme-substrate complexes. Therefore, it was suggested that the changes of cleavage behavior of the modified DNAzymes could be attributed to the influence of the chemical modifications on the cleavage step, instead of the thermal stability of the complex. The cleavage reaction of these DNAzymes on the ³²P-labeled target DNA–RNA–DNA substrate was evaluated under single-turnover conditions.⁴⁵

It has been demonstrated that compound **1** at A9 position of 10–23 DNAzyme contributed a 12-fold increase of cleavage rate, **DZ-1-9** with k_{obs} of 0.0037 min^{−1}⁴⁵. **DZ-1-9** is the fastest DNAzyme in the screening of five dA residues with compound **1** in the catalytic core of 10–23 DNAzyme. Instead of these five dA positions, other residues in the catalytic core were substituted by compound **1** to explore the possible positive role of the extra amino group. Firstly, a position shift of compound **1** around A9 was conducted. When compound **1** in **DZ-1-9** was shifted to its 5'-residue position T8 (**DZ-1-8**), as shown in Fig. 2, **DZ-1-8** could also conduct the reaction like the parent 10–23 DNAzyme, but with a slower cleavage rate (0.0020 ± 0.0002 /min) (Table 2). This replacement resulted in moderate loss of activity, compared with 10–23 DNAzyme, which was not so consistent with previous observation that T8 is the least conserved residue, any other canonical residue or even modified residue replacement results in little effect on the cleavage of 10–23 DNAzyme^{40,41}. On the other hand, the replacement of compound **1** at its 3'-positioned C10 led to a further decrease of the reaction rate, as

Table 1 MALDI-TOF MS data of 10–23 DNAzyme analogs.

DNAzyme	Sequences (5'–3')	MW (calcd.)	MW (found)
DZ-1-8	d(tgc tct cca GGC TAG C1A CAA CGA cct gca cct)	10,060.5	10,062.5
DZ-1-10	d(tgc tct cca GGC TAG CTA 1AA CGA cct gca cct)	10,075.6	10,077.4
DZ-1-13	d(tgc tct cca GGC TAG CTA CAA 1GA cct gca cct)	10,075.6	10,077.0
DZ-1-14	d(tgc tct cca GGC TAG CTA CAA C1A cct gca cct)	10,035.6	10,036.6




Figure 2 The position-dependent effect of compound **1** in the catalytic loop of 10–23 DNAzyme: at A9 (**DZ-1-9**, solid diamond)⁴⁵, at T8 (**DZ-1-8**, solid triangle), at C10 (**DZ-1-10**, open diamond), at C13 (**DZ-1-13**, open square), at G14 (**DZ-1-14**, open triangle), and 10–23 DNAzyme (solid square)⁴⁵.

Table 2 Observed rate constants of 10–23 DNAzyme and its analogs measured under single-turnover conditions.

DNAzyme	T_m (°C)	k_{obs} /min
10–23 DNAzyme	52	0.0037 ± 0.0007 ⁴⁵
DZ-1-9	51	0.045 ± 0.004 ⁴⁵
DZ-1-8	51	0.0020 ± 0.0002
DZ-1-10	51	0.0012 ± 0.0001
DZ-1-13	51	ND
DZ-1-14	51	ND

ND: not detected under present conditions.

observed with **DZ-1-10** ($0.0012 \pm 0.0001 \text{ min}^{-1}$). It meant that the occupation of cytosine at this position was necessary³⁸. Up to now, deletion of C10 or its replacement with other natural or modified residues always resulted in moderate to significant loss of cleavage activity.

The amino groups of C13 and G14 have been demonstrated to be very important for the function of 10–23 DNAzyme³⁸, their amino groups are supposed to be involved in the reaction in some way. When an extra amino group was introduced to these two positions by compound **1**; however, such changes led to a significant loss of the catalytic function, as shown by **DZ-1-13** and **DZ-1-14** (Fig. 2). It seems that the new interactions related to compound **1** at these two positions was detrimental to the reaction, other functional groups in C13 and G14 as well as their spacial occupation might be important for their roles in the context of the catalytic loop.

As demonstrated in our work, in the total nine modified positions (A5, T8, A9, C10, A11, A12, C13, G14 and A15) in the catalytic core of 10–23 DNAzyme⁴⁵, the effect of compound **1** was position-dependent. These observations emphasized the importance of the location for the extra amino group from compound **1** and the conservation of the residues defined by *in vitro* selection. In other words, the catalytic core of 10–23 DNAzyme is very sensitive to chemical modifications, no matter

more or less catalytic power of the new DNAzymes. More importantly, we learnt from these results that A9 is the most appropriate position for the introduction of 7-substituted 8-aza-7-deaza-2'-deoxyadenosine analogs, up to now. This screening research promised more efficient deoxyribozymes for practical applications by chemical functionalization at A9 in the catalytic core of 10–23 DNAzyme.

3. Experimental

3.1. Oligodeoxynucleotides

Oligodeoxynucleotides were synthesized on an ABI 392 DNA/RNA synthesizer (Applied Biosystems, USA) with the DMT-off mode, on a 1 μmol scale according to the User Protocol. The phosphoramidites of canonical residues were purchased from Proligo (Sigma-Aldrich). The oligodeoxynucleotides were deprotected in conc. aq. ammonia for 18 h at 60 °C and purified by a 20% polyacrylamide/7 M urea denaturing gel. After desalting with SEP-PAK cartridges (Oasis MAX, C18, Waters, USA), the product was characterized by MALDI-TOF performed on an autoflex TOF/TOF (BRUKER, Daltonics) with HPA as the matrix. The chimeric substrate was purchased from Takara (Dalin, China). T_m measurement was conducted on a Cary-100 Bio UV-Visible spectrophotometer equipped with a Cary temperature controller (Varian, USA)⁴⁵.

3.2. Cleavage reaction under single-turnover conditions⁴⁵

Prior to the cleavage reaction, the solution of deoxyribozyme (2 μM) and the ³²P-labeled substrate⁴⁵ (20 nM) in 50 mM Tris-HCl (pH 7.4) was denatured at 90 °C for 3 min and cooled to 37 °C. Mg^{2+} was added to a final concentration of 2 mM, and the cleavage reaction was initiated. At different time points, aliquots were taken from the reaction mixture and quenched by an equal volume of stopping solution (100 mM EDTA, 8 M Urea). The substrate and the cleavage product in the samples were separated by electrophoresis on a 20% polyacrylamide gel containing 7 M urea, and quantified by the densitometry of the gel images with a Molecular Dynamics Storm 840 Phosphorimager. The observed rate constant k were calculated according to the equation $P\% = P_{\infty}\% - C \cdot \exp[-k_{\text{obs}}t]$, where $P\%$ is the cleavage percentage of product at time t , $P_{\infty}\%$ is the final percentage of the product at $t = \infty$, C is the difference in $P\%$ between $t = \infty$ and $t = 0$. The data was given as the averaged results of three independent experiments. Less than 20% variation was observed for identical experiments performed on different days.

Acknowledgments

This work was supported by the National Key Technologies R&D Program for New Drugs (2009ZX09301-002) and Natural Science Foundation of China (21072229).

References

1. Santoro SW, Joyce GF. A general purpose RNA-cleaving DNA enzyme. *Proc Natl Acad Sci USA* 1997;**94**:4262–6.
2. Santoro SW, Joyce GF. Mechanism and utility of an RNA-cleaving DNA enzyme. *Biochemistry* 1998;**37**:13330–42.

3. Zhang L, Gasper WJ, Stass SA, Ioffe OB, Davis MA, Mixson AJ. Angiogenic inhibition mediated by a DNAzyme that targets vascular endothelial growth factor receptor 2. *Cancer Res* 2002;**62**:5463–9.
4. Dass CR, Saravolac EG, Li Y, Sun LQ. Cellular uptake, distribution, and stability of 10–23 deoxyribozymes. *Antisense Nucleic Acid Drug Dev* 2002;**12**:289–99.
5. Dass CR. Deoxyribozymes: cleaving a path to clinical trials. *Trends Pharmacol Sci* 2004;**25**:395–7.
6. Schubert S, Kurreck J. Ribozyme- and deoxyribozyme-strategies for medical applications. *Curr Drug Targets* 2004;**5**:667–81.
7. Dass CR. Preclinical anticancer activity of DNA-based cleavage molecules. *Drug Dev Ind Pharm* 2006;**32**:1–5.
8. Dass CR, Choong PFM, Khachigian LM. DNAzyme technology and cancer therapy: cleave and let die. *Mol Cancer Ther* 2008;**7**:243–51.
9. Benson VL, Khachigian LM, Lowe HC. DNAzymes and cardiovascular disease. *Br J Pharmacol* 2008;**154**:741–8.
10. Baum DA, Silverman SK. Deoxyribozymes: useful DNA catalysts *in vitro* and *in vivo*. *Cell Mol Life Sci* 2008;**65**:2156–74.
11. Chan CWS, Khachigian LM. DNAzymes and their therapeutic possibilities. *Int Med J* 2009;**39**:249–51.
12. Unwalla H, Chakraborti S, Sood V, Gupta N, Banerjee AC. Potent inhibition of HIV-1 gene expression and TAT-mediated apoptosis in human T cells by novel mono- and multitarget anti-TAT/Rev/Env ribozymes and a general purpose RNA-cleaving DNA-enzyme. *Antiviral Res* 2006;**72**:134–44.
13. Lu ZX, Ye M, Yan GR, Li Q, Tang M, Lee LM, et al. Effect of EBV LMP1 targeted DNAzymes on cell proliferation and apoptosis. *Cancer Gene Ther* 2005;**12**:647–54.
14. Nunamaker EA, Zhang HY, Shirasawa Y, Benoit JN, Dean DA. Electroporation-mediated delivery of catalytic oligodeoxynucleotides for manipulation of vascular gene expression. *Am J Physiol Heart Circ Physiol* 2003;**285**:2240–7.
15. Xie YY, Zhao XD, Jiang LP, Liu HL, Wang LJ, Fang P, et al. Inhibition of respiratory syncytial virus in cultured cells by nucleocapsid gene targeted deoxyribozyme (DNAzyme). *Antiviral Res* 2006;**71**:31–41.
16. Yu SH, Wang TH, Au LC. Specific repression of mutant K-RAS by 10–23 DNAzyme: sensitizing cancer cell to anti-cancer therapies. *Biochem Biophys Res Commun* 2009;**378**:230–4.
17. Richards JL, Seward GK, Wang YH, Dmochowska IJ. Turning the 10–23 DNAzyme on and off with light. *ChemBiochem* 2010;**11**:320–4.
18. Mokany E, Bone SM, Young PE, Doan TB, Todd AV. MNAszymes a versatile new class of nucleic acid enzymes that can function as biosensors and molecular switches. *J Am Chem Soc* 2010;**132**:1051–9.
19. Hengensbach M, Meusbürger M, Lyko F, Helm M. Use of DNAzymes for site-specific analysis of ribonucleotide modifications. *RNA* 2008;**14**:180–7.
20. Gerasimova YV, Kolpashchikov DM. Nucleic acid detection using MNAszymes. *Chem Biol* 2010;**17**:104–6.
21. Jadhav VM, Scaria V, Maiti S. Antagomirzymes: oligonucleotide enzymes that specifically silence microRNA function. *Angew Chem Int Ed* 2009;**48**:2557–60.
22. Young DD, Lively MO, Deiters A. Activation and deactivation of DNAzyme and antisense function with light for the photochemical regulation of gene expression in mammalian cells. *J Am Chem Soc* 2010;**132**:6183–93.
23. Lusich H, Young DD, Lively MO, Deiters A. Photochemical DNA activation. *Org Lett* 2007;**9**:1903–6.
24. Chen Y, Wang M, Mao C. An autonomous DNA nanomotor powered by a DNA enzyme. *Angew Chem Int Ed* 2004;**43**:3554–7.
25. Schubert S, Gül DC, Grunert HP, Zeichhardt H, Erdmann VA, Kurreck J. RNA cleaving ‘10–23’ DNAzymes with enhanced stability and activity. *Nucleic Acids Res* 2003;**31**:5982–92.
26. Asanuma H, Hayashi H, Zhao J, Liang X, Yamazawa A, Kuramochi T, et al. Enhancement of RNA cleavage activity of 10–23 DNAzyme by covalently introduced intercalator. *Chem Commun* 2006:5062–4.
27. Lam CH, Perrin DM. Introduction of guanidinium-modified deoxyuridine into the substrate binding regions of DNAzyme 10–23 to enhance target affinity: implications for DNAzyme design. *Bioorg Med Chem Lett* 2010;**20**:5119–22.
28. Vester B, Lundberg LB, Sørensen MD, Babu BR, Douthwaite S, Wengel J. LNAzymes: incorporation of LNA-type monomers into DNAzymes markedly increases RNA cleavage. *J Am Chem Soc* 2002;**124**:13682–3.
29. Hollenstein M, Hipolito CJ, Lam CH, Perrin DM. A self-cleaving DNA enzyme modified with amines, guanidines and imidazoles operates independently of divalent metal cations (M^{2+}). *Nucleic Acids Res* 2009;**37**:1638–49.
30. Smuga D, Majchrzak K, Sochacka E, Nawrot B. RNA-cleaving 10–23 deoxyribozyme with a single amino acid-like functionality operates without metal ion cofactors. *New J Chem* 2010;**34**:934–48.
31. Tack F, Bakker A, Maes S, Dekeyser N, Bruining M, Elissen-Roman C, et al. Modified poly(propylene imine) dendrimers as effective transfection agents for catalytic DNA enzymes (DNAzymes). *J Drug Target* 2006;**14**:69–86.
32. Kubo T, Takamori K, Kanno K, Bakalova R, Ohba H, Matsukisano M, et al. Efficient cleavage of RNA, enhanced cellular uptake, and controlled intracellular localization of conjugate DNAzymes. *Bioorg Med Chem Lett* 2005;**15**:167–70.
33. Okumoto Y, Tanabe Y, Sugimoto N. Factors that contribute to efficient catalytic activity of a small Ca^{2+} -dependent deoxyribozyme in relation to its RNA cleavage function. *Biochemistry* 2003;**42**:2158–65.
34. Ota N, Warashina M, Hirano K, Hatanaka K, Taira K. Effects of helical structures formed by the binding arms of DNAzymes and their substrates on catalytic activity. *Nucleic Acids Res* 1998;**26**:3385–91.
35. Perrin DM, Garestier T, Hélène C. Briding the gap between proteins and nucleic acids: a metal-independent RNaseA mimic with two protein-like functionalities. *J Am Chem Soc* 2001;**123**:1556–63.
36. Sidorov AV, Grasby JA, Williams DM. Sequence-specific cleavage of RNA in the absence of divalent metal ions by a DNAzyme incorporating imidazolyl and amino functionalities. *Nucleic Acids Res* 2004;**32**:1591–601.
37. Thomas JM, Yoon JK, Perrin DM. Investigation of the catalytic mechanism of a synthetic DNAzyme with protein-like functionality: an RNaseA mimic? *J Am Chem Soc* 2009;**131**:5648–58.
38. Nowakowski J, Shim PJ, Prasad GS, Stout CD, Joyce GF. Crystal structure of an 82-nucleotide RNA-DNA complex formed by the 10–23 DNA enzyme. *Nat Struct Biol* 1999;**6**:151–6.
39. Kenward M, Dorfman KD. Coarse-grained brownian dynamics simulations of the 10–23 DNAzyme. *Biophys J* 2009;**97**:2785–93.
40. Zaborowska Z, Fürste JP, Erdmann VA, Kurreck J. Sequence requirements in the catalytic core of the 10–23 DNA enzyme. *J Biol Chem* 2002;**277**:40617–22.
41. Zaborowska Z, Schubert S, Kurreck J, Erdmann VA. Deletion analysis in the catalytic region of the 10–23 DNA enzyme. *FEBS Lett* 2005;**579**:554–8.
42. Nawrot B, Wiedera K, Wojcik M, Rebowska B, Nowak G, Stec WJ. Mapping of the functional phosphate groups in the catalytic core of deoxyribozyme 10–23. *FEBS J* 2007;**274**:1062–72.
43. Wang B, Cao L, Chiuman W, Li Y, Xi Z. Probing the function of nucleotides in the catalytic cores of the 8–17 and 10–23 DNAzymes by abasic nucleotide and C3 spacer substitutions. *Biochemistry* 2010;**49**:7553–62.
44. Robaldo L, Montserrat JM, Iribarren AM. 10–23 DNAzyme modified with (2' R)- and (2' S)-2'-deoxy-2'-C-methyluridine in the catalytic core. *Bioorg Med Chem Lett* 2010;**20**:4367–70.
45. He JL, Zhang D, Wang Q, Wei X, Cheng MS, Liu KL. A novel strategy of chemical modification for rate enhancement of 10–23 DNAzyme: a combination of A9 position and 8-aza-7-deaza-2'-deoxyadenosine analogs. *Org Biomol Chem* 2011;**9**:5728–36.